

Increase in glycaemia stimulates reactive oxygen species (ROS) production by polymorphonuclear neutrophils in type 2 diabetic patients

Dorota Pisarczyk-Wiza¹, Dorota Zozulińska-Ziółkiewicz¹, Henryk Wysocki²,
Bogna Wierusz-Wysocka¹

¹ Department of Internal Medicine and Diabetology, University of Medical Sciences, Poznan, Poland

² Department of Cardiology and Intensive Therapy, University of Medical Sciences, Poznan, Poland

Abstract

Polymorphonuclear neutrophils (PMNs), oxidative stress and hyperglycaemia play important roles in the development of micro- and macroangiopathy in diabetes. The aim of this study was to evaluate the influence of a meal and the increment of postprandial glycaemia on reactive oxygen species production by PMNs in type 2 diabetic patients.

The study was performed on 25 patients – 18 women and 7 men, aged 61.7±10.5 years, diagnosed with type 2 diabetes for 16.0±7.5 years, and in 20 healthy control subjects. PMNs were isolated from the blood by single-step gradient centrifugation. The superoxide anion (O₂⁻) production by PMNs was measured spectrophotometrically by cytochrome C reduction, and the hydrogen peroxide (H₂O₂) level was measured spectrophotometrically by phenyl red oxidation. A significant correlation was observed between an increase in glycaemia and O₂⁻ production (r=0.52, p<0.05) and in H₂O₂ production by PMNs (r=0.43, p<0.05). The results obtained suggest that in obese type 2 diabetic patients it is the increment of postprandial glycaemia, and not directly the meal itself, that stimulates reactive oxygen species production by PMNs. This might play an important role in the development of late vascular complications and thus have clinical implications.

Key words

reactive oxygen species, polymorphonuclear neutrophils, postprandial state, diabetes mellitus type 2

INTRODUCTION

Chronic complications of diabetes continue to create a significant clinical problem. Macroangiopathy, microangiopathy and diabetic neuropathy lead to severe disability and premature death. To date, the responsible pathogenetic mechanisms have not been fully understood; nevertheless, hyperglycaemia is known to involve a factor which plays a key role in the induction and progression of chronic diabetic complications. This was proved in the Diabetes Control and Complications Trials (DCCT) studies performed on type 1 diabetic patients, as well as in both the Kumamoto Study and the United Kingdom Prospective Diabetes Study (UKPDS) which involved patients with type 2 diabetes [1-3]. All these studies clearly demonstrated that efforts to reach normoglycaemia and optimal metabolic control significantly reduce the complications of diabetes, especially those involving microangiopathy. In recent years attention has been drawn to the role played by abnormal postprandial glycaemia. Its role in the development of atherosclerotic lesions was observed by Hanefeld et al. [4]. Their results indicate that postprandial hyperglycaemia represents an independent risk factor of death in patients with newly diagnosed diabetes. In addition, several other, earlier reports, including the Honolulu Heart Study,

Whitehall Study, Paris Prospective Study and the Helsinki Policemen's Study, all showed that even a slight increase in the blood glucose level following a meal significantly increased the risk of cardiovascular disease [5-7]. Ceriello et al. have suggested that postprandial increases in the blood glucose concentration may lead to abnormalities in the function and structure of cells in the vascular wall [8]. Together with others, Ceriello et al. noticed that hyperglycaemia may be responsible for the pronounced apoptosis of these cells due to intensified protein glycation, increased production of reactive oxygen species, and alterations in the clotting and fibrinolysis mechanisms through the activation of protein kinase C, and of the transcription nuclear factor NFκ-B [8-10].

In recent years, the inflammatory theory of the development of chronic diabetic complications has gained an increasing number of adherents [11-13]. The syndrome of hyperglycaemia-induced disturbances corresponds to lesions which can be detected at inflammatory foci. The principal cells involved in a non-specific inflammatory response are polymorphonuclear neutrophils (PMNs). These circulate in the blood for only 6-8 hours at rest, but in a state of readiness they become stimulated. If not stimulated, i.e. in normal conditions, a significant proportion of PMNs undergo apoptosis. The remaining fraction leaves the circulation and undergoes destruction in the tissues, or on the surface of mucous membranes. Activated PMNs alter their properties and adjust to chemotaxis and to interaction with themselves (aggregation) and to other cells, such as monocytes, blood platelets, endothelial cells (adhesion). The PMN activation stage involves diapedesis and migration of cells through the

Corresponding author: Dorota Pisarczyk-Wiza, Department of Internal Medicine and Diabetology, University of Medical Sciences, Raszeja Hospital, Mickiewicza 2, 60-834 Poznan, Poland; tel.: +4861 844579; fax: +48618474579.
E-mail: wizus@wp.pl

Received: 13 April 2011; accepted: 27 June 2011

endothelium. They reach the sites at which inflammation-inducing factors are acting and there phagocytose the foreign bodies. The process of degradation of phagocytised material involves the reactive oxygen species (ROS) and proteolytic enzymes released into the phagosome. The condition in which increased activities of reactive oxygen species are detected used to be termed 'oxidative stress' [14,15]. The stress develops due to disturbed equilibrium between ROS production and the capacity to eliminate them by anti-oxidative systems. The reactive oxygen species include, among others, superoxide anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), hydroxyl radical ($\cdot OH$), hydrogen superoxide (HO_2), single oxygen (1O_2) and nitric oxide (NO) [14,16,17]. ROS may manifest their toxic activity not only within a phagosome but also outside the cell. The phenomenon develops in the presence of chemotactic factors. Mediators of the inflammatory response, released from activated PMNs, induce destructive processes which affect the neighbouring tissues.

Apart from the action of classical inflammatory agents, activation of PMN may also result from the action of metabolic factors. In conditions of hyperglycaemia the cells manifest permanent activation and therefore a reduced response to standard stimulatory agents [18-20]. It was suggested recently that the intensity of the atherogenic inflammatory process may reflect, among other factors, superphysiological increments in glycaemia. We therefore decided to determine the effect of a meal on reactive oxygen species release from PMNs.

MATERIALS AND METHODS

Evaluation of the influence of ingesting a standard meal on ROS production by PMNs was performed on 25 patients with type 2 diabetes – including 18 women and 7 men, 44-74 years of age, with a history of clinically evident diabetes for periods ranging from 2 months to 30 years. All the diabetic patients met the criteria for the diagnosis of the disease: i.e. they manifested the abdominal type of obesity, decreased sensitivity of peripheral tissues to insulin, arterial hypertension and hyperlipidaemia.

The control group consisted of 20 healthy individuals, 10 women and 10 men, 30.0 ± 8.9 years of age and with a BMI of 22.5 ± 1.2 kg/m². In this group, blood pressure levels did not exceed 130/80 mmHg and the values of glycaemia and the lipid parameters fitted the appropriate normal range, and the serum C-reactive protein level was less than 3 mg/l.

Patients with renal and/or hepatic failure, persons with acute inflammatory conditions and tobacco smokers were excluded from the study.

All the participants were informed about the purpose of the investigation and gave their written consent. The schedule of this study (within a broader research project on the role of the inflammatory process, with particular attention devoted to PMNs, in the development of chronic complications in diabetes) was approved by the Ethics Committee of Poznan University of Medical Sciences.

The clinical characteristics of the subjects of this investigation are presented in Table 1.

The parameters of metabolic control were determined in all the patients, using HOMA (Homeostasis Model Assessment), the sensitivity of peripheral tissues to insulin (HOMA-IR) and the function of pancreatic beta cells (HOMA-B) were calculated as shown below:

Table 1. Clinical characteristics of patients with type 2 diabetes and controls

	25	20
Number (n)	25	20
Gender (females/males)	18/7	10/10
Age (years)	61.7 ± 10.5	30.0 ± 8.9
Diabetes duration (years)	16.0 ± 7.5	-
Fasting plasma glucose (mg/dl) [mmol/l]	157.4±51.5 [8.7±2.8]	70.3±10.1 [3.8±0.6]
Glycaemia 2 hours after a meal (mg/dl) [mmol/l]	229.5±77.9 [12.7±4.3]	74.4±8.0 [4.1±0.4]
Increment of glycaemia (mg/dl) [mmol/l]	72.1±64.7 [4.0±3.9]	-
HbA1c (%)	8.4±0.9	5.1±0.8
Fructosamine (µmol/l)	385.6±67.8	187.2±27.8
Total cholesterol (mg/dl) [mmol/l]	232.2±38.0 [6.0±1.0]	181.7±34.8 [4.7±0.9]
HDL-cholesterol (mg/dl) [mmol/l]	41.6±19.7 [1.1±0.5]	46.4±11.6 [1.2±0.3]
LDL-cholesterol (mg/dl) [mmol/l]	142.6±39.0 [3.7±1.0]	92.8±11.6 [2.4±0.3]
Triglycerides (mg/dl) [mmol/l]	267.7±170.8 [3.0±1.9]	106.3±35.4 [1.2±0.4]
BMI (kg/m ²)	34.9±6.3	22.5±1.2
WHR	1.0±0.1	0.9±0.1
Systolic BP (mmHg)	146±14	116±18
Diastolic BP (mmHg)	85±10	68±10
WBC level upon fasting (x10 ⁹ /l)	5.6±1.4	6.3±0.4
WBC level 2 hours after a meal (x10 ⁹ /l)	6.0±1.9	7.1±0.6
CRP (mg/l)	6.4±4.8	1.1±0.1
HOMA-IR (%)	13.0±2.8	-
HOMA-B (%)	158.6±29.9	-
Microalbuminuria (n)	11	-
Retinopathy (n)	17	-
Neuropathy (n)	5	-
Stroke (n)	3	-
Ischaemic cardiac disease (n)	10	-
Insulin alone (n)	12	-
Insulin + metformin (n)	6	-
Metformin (n)	3	-
Metformin + Sulfonylurea (n)	2	-
Diet only (n)	2	-
ACE-inhibitors (n)	25	-
Statins (n)	10	-
Fibrates (n)	5	-
Acetylosalicylic acid (n)	11	-
Indapamide (n)	6	-
Calcium channel antagonists (n)	7	-
Beta-blockers (n)	8	-

HbA1c – glycated haemoglobin; HDL – high density lipoproteins; LDL – low density lipoproteins; BMI – body mass index; WBC – white blood cells; CRP – C-reactive protein; HOMA-IR – Homeostasis Model Assessment-insulin resistance; HOMA-B – Homeostasis Model Assessment-beta cells function.

HOMA-IR = insulinaemia upon fasting (µU/ml) × glycaemia upon fasting (mmol/l) / 22.5.

HOMA-B = insulinaemia upon fasting (µU/ml) × 20 / [glycaemia upon fasting (mmol/l) – 3.5].

Studies were performed in all the diabetic patients to detect any chronic complications of the disease. Retinopathy was diagnosed by examination of the fundus of the eyes after

dilatation of the pupils. Nephropathy was diagnosed by the presence of microalbuminuria (2 positive results of 3 tests). The presence of any neuropathy was diagnosed on the basis of the history, tests of the Achilles tendon reflex, sense of touch and vibration. The presence of cardiovascular disease was detected via the history, 12-lead ECG examination, echocardiography and evaluation of the pulse in the dorsal arteries of the feet.

In all subjects the parameters of metabolic control (fasting plasma glucose level, postprandial plasma glucose level, serum fructosamine level, serum concentration of total cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides) were evaluated using a standard technique and the HbA1c level by means of high liquid chromatography (HPLC).

The white blood cell (WBC) count in the fasting state, and 2 hours after a meal, were assessed using standard laboratory techniques. The serum insulin concentration was measured using the immunoenzymatic technique, and C-reactive protein by the highly sensitive immunoturbidimetric technique using HITACHI 717 kits.

For studies on the function of PMNs, 6 ml of peripheral blood was taken from a cubital vein into plastic tubes containing heparin to a final concentration of 1 unit/ml blood. The blood was sampled twice: before and 2 hours after a standard meal rich in carbohydrates, containing 30 g (126 kcal or 60%) carbohydrates, 6g (54 kcal or 25%) fat and 8g (34 kcal or 15%) protein. The total calorific value of the meal amounted to approximately 214 kcal. The studies were performed on PMNs isolated from the venous blood.

Isolation of PMNs according to a modified version of the Boyum technique [21]. Six ml of blood was overlaid on 3 ml of Gradisol G preparation and centrifuged for 30 min at 2,000 rpm. After collection of the PMN-rich buffy coat, the cells were washed twice with buffered saline. The remaining admixture of erythrocytes in the sediment was lysed using a hypotonic lysing agent (a reagent containing 0.828 g NH_4Cl , 0.1 g KHCO_3 dissolved in 100 ml distilled water, with the addition of 20 μl 1 N Na OH). The isolated PMN were suspended in a modified Hanks solution (MHS) (11mM glucose, 9 mM K_2HPO_4 , 145 mM NaCl, 1 mM HCl, pH 7.4), in the volume necessary to obtain a final cell concentration appropriate for the planned studies 5×10^6 cells/ml. Using this technique it was possible to recover 70%-90% of the original cell content. The final suspension contained around 98% PMNs. PMN numbers were scored in a Bürker's haemocytometer, following their dilution in Türk's reagent.

The percentage of dead cells, determined following every isolation using 0.5% trypan blue, did not exceed 6%. Studies on PMN function were completed before 3 hours had elapsed since the time when the suspension of the cells had been obtained.

Production of superoxide anions ($\text{O}_2^{\cdot-}$) by PMNs, according to the modified technique of Babior et al. [22]. All 96 wells in a microtitration plate were charged with 50 μl of the PMN suspension (5×10^6 cells/ml) and with 50 μl of cytochrome C solution (5 mg/ml, Sigma). Subsequently, 50 μl MHS was added to each well. The blind test was supplemented with 50 μl superoxide dismutase (SOD) at a concentration of 30 U/ml. The plate thus prepared was incubated for 30 min at a temperature of 37°C. Subsequently, 50 μl SOD solution was added to the samples and the plate was evaluated using

spectrophotometry and an ELISA reader at the wavelength of $\lambda=550$ nm. The results were expressed in $\text{nmol}/5 \times 10^6 \text{PMNs}/30$ min.

Production of hydrogen peroxide (H_2O_2) by PMN according to the modified technique of Pick [23]. Each of the 96 wells in a microtitration plate were charged with 25 μl of the suspension of the PMN (5×10^6 cells/ml) and with 50 μl of phenol red solution. Subsequently, 25 μl MHS was added to each well. Following incubation (30 min., temp. 37°C), the reaction was stopped by the addition of 10 μl 1N NaOH. The amount of H_2O_2 was evaluated spectrophotometrically at the wave length of $\lambda=600$ nm, using a standard curve, drawn every time for a row of respective hydrogen peroxide dilutions. The results were expressed in $\text{nmol}/5 \times 10^6 \text{PMNs}/30$ min.

Statistical evaluation of the results. Normality of distribution was tested using the Kolmogorov-Smirnov test.

In cases of normal distribution in the results of the effect of a meal on PMN function, Student's *t* test was used. When the results manifested any distribution that differed from the normal, the Wilcoxon's non-parametric test was used.

Differences between the groups being evaluated were appraised using the Mann-Whitney test. The relationship between increases in glycaemia and reactive oxygen species production by PMNs was assessed using the Pearson (*r*) correlation test. Values at $p < 0.05$ were accepted as representing significant differences.

The results of $\text{O}_2^{\cdot-}$ and H_2O_2 production by PMNs were presented as means \pm standard error of the mean (SEM).

Statistical computations were facilitated by application of GraphPadInstat w.3.05 software.

RESULTS

In the group of healthy individuals, fasting glycaemia averaged $70.3 \pm 10.1 \text{mg/dl}$ ($3.8 \pm 0.6 \text{mmol/l}$); 2 hours after the meal it was $74.4 \pm 8.0 \text{mg/dl}$ ($4.1 \pm 0.4 \text{mmol/l}$). No significant postprandial increase in glycemia was found ($p > 0.05$).

In the patients with type 2 diabetes, fasting glycaemia averaged $157.4 \pm 51.5 \text{mg/dl}$ ($8.7 \pm 2.8 \text{mmol/l}$), this being significantly higher than in healthy individuals ($p < 0.05$). Two hours after the meal, glycaemia increased significantly to a mean of $229.5 \pm 77.9 \text{mg/dl}$ ($12.7 \pm 4.3 \text{mmol/l}$) ($p < 0.05$). In all the type 2 diabetic patients the meal induced an increase in glycaemia by a mean of $72.1 \pm 64.7 \text{mg/dl}$ ($4.0 \pm 3.9 \text{mmol/l}$).

In the healthy subjects, the mean WBC level before the standard meal was $6.3 \pm 0.4 (\times 10^9/l)$, and following the meal it rose to $7.1 \pm 0.6 (\times 10^9/l)$. In the type 2 diabetic patients, mean WBC values were similar being, respectively, before the meal: $5.6 \pm 1.4 (\times 10^9/l)$ and following the meal $6.0 \pm 1.9 (\times 10^9/l)$.

Effect of the test meal and the postprandial increase in glycaemia on the production of superoxide anions ($\text{O}_2^{\cdot-}$) by PMNs (Figs. 1 and 2). Production of $\text{O}_2^{\cdot-}$ upon fasting was significantly higher in the type 2 diabetic patients than in the healthy individuals ($8.40 \pm 1.11 \text{nmol}/5 \times 10^6 \text{PMNs}/30 \text{min}$ vs. $4.28 \pm 0.59 \text{nmol}/5 \times 10^6 \text{PMN}/30 \text{min}$, $p < 0.01$). On the other hand, 2 hours after the meal, no significant differences in $\text{O}_2^{\cdot-}$ production could be detected between the 2 groups




Figure 1. Production of O_2^- by isolated PMNs

* $p < 0.05$ before vs after the meal in healthy individuals,
 $\Delta p < 0.05$ in type 2 diabetic patients vs healthy individuals upon fasting.




Figure 2. Relationship between postprandial increment in glycaemia and production of O_2^- by PMNs 2 hours after a meal in type 2 diabetic patients

(10.54 ± 2.65 nmol/ 5×10^6 PMN/30min vs. 7.10 ± 1.14 nmol/ 5×10^6 PMNs/30min, $p > 0.05$).

In the healthy individuals, a significant increase in O_2^- production was detected 2 hours after the standard meal (4.28 ± 0.59 nmol/ 5×10^6 PMNs/30min. vs. 10.54 ± 2.65 nmol/ 5×10^6 PMN/30min, $p < 0.05$).

The meal did not significantly change the production of O_2^- in type 2 diabetic patients (8.40 ± 1.11 nmol/ 5×10^6 PMNs/30min vs. 7.10 ± 1.14 nmol/ 5×10^6 PMNs/30min, $p > 0.05$). However, in the diabetic patients, the production of O_2^- by PMNs in the second hour after the meal demonstrated a significant correlation with the postprandial increase in glycaemia ($r = 0.52$, $p < 0.05$). No correlation was found between the postprandial increase in glycaemia and O_2^- production by PMNs in the healthy controls ($p > 0.05$).

Effect of the test meal and postprandial increase in glycaemia on the production of hydrogen peroxide (H_2O_2) by PMNs (Figs. 3 and 4). No significant differences were discovered between the examined groups in the production of H_2O_2 by PMNs upon fasting (20.68 ± 1.20 nmol/ 5×10^6 PMNs/30min vs. 24.01 ± 1.68 nmol/ 5×10^6 PMNs/30min, $p > 0.05$) or 2 hours after the standard meal (21.99 ± 1.24 nmol/ 5×10^6 PMNs/30min vs. 32.57 ± 3.76 nmol/ 5×10^6 PMNs/30min, $p > 0.05$).

Two hours after the meal, in comparison to the fasting state, significantly higher H_2O_2 production was detected in the group of healthy individuals (24.01 ± 1.68 nmol/ 5×10^6 PMN/30min. vs. 32.57 ± 3.76 nmol/ 5×10^6 PMN/30min, $p < 0.05$).

The meal failed to significantly affect the production of hydrogen peroxide by PMNs in patients with type 2 diabetes (20.68 ± 1.20 nmol/ 5×10^6 PMNs/30min vs. 21.99 ± 1.24 nmol/ 5×10^6 PMNs/30min, $p > 0.05$). In these patients, the production of H_2O_2 by PMNs 2 hours after the meal was found to exhibit




Figure 3. Production of H_2O_2 by isolated PMNs

* $p < 0.05$ before vs after the meal, in healthy individuals




Figure 4. Relationship between postprandial increment in glycaemia and production of H_2O_2 by PMN 2 hours after the meal in type 2 diabetic patients

a significant correlation with the postprandial increment in glycaemia ($r = 0.44$, $p < 0.05$). No such correlation was found between the postprandial increase in glycaemia and H_2O_2 production by PMNs in the healthy controls ($p > 0.05$).

DISCUSSION

Polymorphonuclear neutrophils appear to play a significant role in the development of diabetic micro- and macroangiopathy. As cells of the inflammatory response they are engaged in processes induced by hyperglycaemia, leading to injury of the vascular endothelium. In diabetic patients, several disturbances of PMN function were noticed [24-27]. Wierusz-Wysocka et al., demonstrated negative effects of increasing glucose concentrations on some PMN functions in *in vitro* conditions [19,20]. PMNs are a significant source of reactive oxygen species, providing basic elements for the oxygen-dependent bactericidal mechanisms. Disturbance in the properties (e.g., an inborn deficit in NADPH oxidase) may lead to severe, frequently lethal infections [28]. An increase in ROS production by PMNs may result from humoral factors, including an acute or chronic hyperglycaemia [29]. Most reports in the literature draw attention to increases in the production of reactive oxygen species by unstimulated PMN and the less effective response to activating factors in diabetic patients [30]. This has been confirmed by the results of our study which showed that the production of O_2^- upon fasting was significantly higher in patients with type 2 diabetes than in normal individuals. Ceriello et al. observed a relationship between the development of oxidative stress and rapid, postprandial increments in glycaemia. These processes include, among others, the stimulated glycation of proteins, auto-oxidation of glucose, and the intracellular activation of glucose turnover along the polyolic pathway [8,9]. In our

study, glycaemia increased following the meal, but no parallel increase was observed in the production of superoxide anions or hydrogen peroxide by PMNs isolated from the blood of patients with type 2 diabetes. The absence of the effects of a meal on the production of ROS by PMNs in diabetics may point to a disturbed reactivity of the cells to additional stimulatory factors. In patients with type 2 diabetes, PMNs and, most probably, their precursors in the bone marrow, are exposed in a chronic fashion to elevated glucose concentrations. Poor metabolic control was manifested by the abnormal, elevated levels of fructosamine and HbA1c at the beginning of the studies. Zozulińska et al. have already shown that PMN function is dependent on the HbA1c value [31]. The additional stimulatory factors might also have involved the hyperlipemia and hypertriglyceridemia, in particular, which we also noted in our studies. Under such conditions, the PMNs probably remain in a permanent state of sub-threshold stimulation. Several authors have suggested that PMNs in this state are unable to respond appropriately to the action of typical stimulatory factors [25,32-34]. The standard meal applied in our studies apparently failed to provide a sufficiently strong stimulus to activate the sub-threshold activated cells. However, attention should be drawn to the significant relationship noted in our studies between the postprandial increment in glycaemia and the production of $O_2^{\cdot-}$ and H_2O_2 by PMNs in patients with type 2 diabetes 2 hours after the meal. The relationship demonstrated is consistent with the suggestions of Ceriello et al., that the augmented ROS production is primarily responsible for the postprandial 'toxicity' of hyperglycaemia (35). Our studies additionally draw attention to the potential role of PMNs in the initiation of the acute process of postprandial damage to vascular walls.

In contrast to the diabetic patients, in healthy individuals the meal exerted a significant effect on the intensity of $O_2^{\cdot-}$ and H_2O_2 production by isolated PMNs. These results are consistent with the work of Mohanty et al., who showed that loading with 75g glucose was followed in the subsequent 3 hours by increased production of ROS by PMNs and monocytes isolated from the blood of healthy individuals [29]. The most pronounced production of reactive oxygen species was recorded in the second hour of the test. They demonstrated an increased release of ROS by PMNs and monocytes obtained from healthy individuals following a meal rich in lipids and proteins [36]. Their results, as well as those obtained in our study, suggest that the function of PMNs in healthy individuals may be dependent on the composition of meals, including the carbohydrate content. The response of PMN to alimentary stimulus shows how cells involved in the inflammatory response behave under physiological conditions. PMNs released from the marginal pool under the effect of a meal remain in the pre-activation phase. In this state, they are characterized by high sensitivity to the action of sequential stimulatory factors and are capable of releasing reactive oxygen species. In the PMNs of healthy individuals, the augmented production of ROS follows the intensity of the intracellular glucose metabolism. Loading with glucose directly activates NADPH oxidase in PMN, the key enzyme for the production of superoxide anions. In turn, the superoxide anion provides a substrate for the cascade generation of ROS [14,15,37]. In physiological conditions, reactive oxygen species exert no particularly destructive effects on tissues: their damaging affect on cells is

counteracted by effective anti-oxidative systems. Ceriello et al. demonstrated that in the course of an oral glucose tolerance test performed on healthy individuals, the increment in blood glucose concentration is paralleled by an increase in blood malonyldialdehyde (MDA) concentration, and by a decrease in the total anti-oxidative capacity of plasma. Apparently, in patients with metabolically unbalanced diabetes, anti-oxidative systems exhibit low efficacy.

The role of PMNs in the development of atherosclerotic lesions seems to be increasingly well documented. Postprandial hyperglycaemia and hyperlipaemia modulate the course of the inflammatory response, including the functions of PMNs. We found that a meal rich in carbohydrates modulates the functions of the PMN cells, and the effect in patients with type 2 diabetes was different from that in normal individuals. Determination of the role of various other factors on the functions of PMNs requires further study.

ABBREVIATIONS

1O_2	– single oxygen
BMI	– Body Mass Index
BP	– Blood Pressure
CRP	– C-reactive protein
DCCT	– Diabetes Control and Complications Trials
H_2O_2	– hydrogen peroxide
HbA1c	– glycated haemoglobin
HDL	– high density lipoproteins
HMPS	– hexomonophosphatate pathway
HO_2	– hydrogen superoxide
HOMA	– Homeostasis model assessment
HOMA-B	– Homeostasis Model Assessment-beta cells function
HOMA-IR	– Homeostasis Model Assessment-insulin resistance
LDL	– low density lipoproteins
MDA	– malonyldialdehyde
MHS	– modified Hanks solution
NFκ-B	– transcription nuclear factor kappa-B
NO	– nitric oxide
$O_2^{\cdot-}$	– superoxide anion
OH	– hydroxy radicals
PKC	– protein kinase C
PMN	– polymorphonuclear neutrophils
ROS	– reactive oxygen species
SOD	– superoxide dismutase
TG	– triglycerides
UKPDS	– United Kingdom Prospective Diabetes Study
WBC	– white blood cell
WHR	– Waist to Hip Ratio

AUTHORSHIP RESPONSIBILITY

Dorota Pisarczyk-Wiza (DPW) contributed to the performance of the study and writing article. Dorota Zozulinska-Ziolkiewicz (DZZ) and Bogna Wierusz-Wysocka (BWW) contributed to the conception and design of the study. Dorota Zozulinska-Ziolkiewicz additionally carried out the statistical analysis of the data. Henryk Wysocki (HW) cooperated in the performance of the study.

ACKNOWLEDGEMENTS

The authors express their thanks to Dr A. Szczepanik and Mgr A. Banaszak in the laboratory of the Department of Cardiology and Intensive Therapy at the University of Medical Sciences in Poznan for their excellent technical assistance and cooperation. The authors are also indebted to Prof. Geoffrey Shaw for his editorial assistance.

REFERENCES

1. United Kingdom Prospective Diabetes Study Group (UKPDS): Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;12:837-53.
2. Shichiri M, Kishikawa H, Ohkubo Y, Wake N. Long-term results of the Kumamoto Study on optimal diabetes control in type 2 diabetes patients. *Diabetes Care* 2000;23(Suppl 2):B21-B29.
3. The Diabetes Control and Complications Trials (DCCT) Research Group: The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *New Engl J Med* 1993;329(14):977-86.
4. Hanefeld M, Fisher S, Julius U, Schulze J, Schwanebeck U, Schmechel H, Zeigelasch HJ, Lindner J. Risk factors for myocardial infarction and death in newly detected NIDDM: the Diabetes Intervention Study, 11-year follow-up. *Diabetologia* 1996;39(12):1577-1583.
5. Danahue R, Abbott R, Reed D. Post challenge glucose concentration and coronary heart disease in men of Japanese ancestry. Honolulu Heart Program. *Diabetes* 1987;36(6):689-692.
6. Balkau B, Shipley M, Jarrett RJ, Pyorala K, Pyorala M, Forhan A, Eschwege E. High blood glucose concentration is a risk factor for mortality in middle-aged non-diabetic men. 20-year follow-up in the Whitehall Study, the Paris Prospective Study and the Helsinki Policemen Study. *Diabetes Care* 1998;21(3):360-367.
7. Fuller JH, Shipley MJ, Rose G, Jarrett RJ, Keen H. Coronary heart disease risk and impaired glucose tolerance: the Whitehall Study. *Lancet* 1980;1(8183):1373-1376.
8. Ceriello A. The emerging role of postprandial hyperglycemic spikes in the pathogenesis of diabetic complications. *Diab Med* 1998;15:188-193.
9. Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001;414:813-820.
10. Ishii H, Koya D, King GL. Protein kinase C activation and its role in the development of vascular complications in diabetes mellitus. *J Mol Med* 1998;76:1-31.
11. Schmid MI, Duncan BB, Sharett AR. Markers of inflammation and prediction of diabetes mellitus in adults (Atherosclerosis risk in communities study): a cohort study. *Lancet* 1999;353:1649-1652.
12. Zozulińska D, Majchrzak A, Markiewicz S, Wierusz-Wysocka B. Selected markers of inflammation in diabetic patients. *Diabetes Res* 1996;3:33-39.
13. Zozulińska D, Wierusz-Wysocka B. Hyperglycemia and inflammation are culprits of late complications. *Arch Med Sci* 2005;1:115-118.
14. Halliwell B. The role of oxygen radicals in human disease, with particular reference to the vascular system. *Haemostasis* 1993;23(Suppl 1):118-126.
15. Freeman BA, Crapo JD. Biology of disease: Free radicals and tissue injury. *Lab Invest* 1982; 47:412-46.
16. Giugliano D, Ceriello A, Paolisso G. Oxidative stress and diabetic vascular complications. *Diabetes Care* 1996;33:257-267.
17. Baynes JW, Thorpe SR. Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *Diabetes* 1999;48(1):1-9.
18. Takahashi T, Hato F, Yamane T, Inaba M, Okuno Y, Nishizawa Y, Kitagawa S. Increased spontaneous adherence of neutrophils from type 2 diabetic patients with overt proteinuria. *Diabetes Care* 2000;23:417-418.
19. Wierusz-Wysocka B, Wysocki H, Byks H, Zozulińska D, Wykrętowicz A, Kaźmierczak M. Metabolic control and free radical activity in diabetic patients. *Diabetes Res Clin Pract* 1995;27(3):193-197.
20. Wierusz-Wysocka B, Wysocki H, Wykrętowicz A, Klimas R. The influence of increasing glucose concentration on selected functions of polymorphonuclear neutrophils. *Acta Diabetol Lat* 1988;25:283-288.
21. Boyum A. Isolation of lymphocytes, granulocytes and macrophages. *Scand J Immunol* 1976;Suppl 5:9-15.
22. Babior BM, Kipnes RS, Curnutte JT. Biological defense mechanisms. The production by leukocytes of superoxide, a potential bactericidal agent. *J Clin Invest* 1973;52(3):741-744.
23. Pick E. Microassays for superoxide and hydrogen peroxide production and nitroblue tetrazolium reduction using an enzyme immunoassay microplate reader. *Methods Enzymol* 1986;132:407-421.
24. Caimi G, Canino B, Montana M, Ventimiglia G, Catania A, Lo Presti R. Polymorphonuclear leukocyte membrane fluidity and cytosolic Ca²⁺ concentration in diabetes mellitus. *Acta Diabetol* 1998;35:158-160.
25. Delamaire M, Maugeudre D, Moreno M, Le Goff MC, Allanic H, Genetet B. Impaired leukocyte functions in diabetic patients. *Diabet Med* 1997;14(1):29-34.
26. Marhoffer W, Stein M, Maeser E, Federlin K. Impairment of polymorphonuclear leukocyte function and metabolic control of diabetes. *Diabetes Care* 1992;15:256-260.
27. Omori K, Ohira T, Uchida Y, Aylavarapu S, Batista EL Jr, Yagi M, Iwata T, Liu H, Hasturk H, Kantarci A, Van Dyke TE. Priming of neutrophil oxidative burst in diabetes requires preassembly of the NADPH oxidase. *J Leukoc Biol* 2008;84:292-301.
28. Morel F. Molecular aspects of chronic granulomatous disease - 'the NADPH oxidase complex'. *Bull Acad Natl Med* 2007;191:377-390.
29. Mohanty P, Hamouda W, Garg R, Aljada A, Ghanim H, Dandona P. Glucose challenge stimulates reactive oxygen species (ROS) generation by leukocytes. *J Clin Endocrinol Metab* 2000;85:2970-2973.
30. Wierusz-Wysocka B, Wysocki H, Siekierka H, Wykrętowicz A, Szczepanik A, Klimas R. The evidence of polymorphonuclear neutrophils (PMN) activation in patients with insulin-dependent diabetes mellitus. *J Leukoc Biol* 1987;42:519-523.
31. Zozulińska D, Wierusz-Wysocka B, Wysocki H, Majchrzak A, Wykrętowicz A. The influence of IDDM duration on superoxide anions and hydrogen peroxide production by polymorphonuclear neutrophils. *Diabetes Res Clin Pract* 1996;33:139-144.
32. Zozulińska D, Majchrzak A, Markiewicz S, Wierusz-Wysocka B. Selected markers of inflammation in diabetic patients. *Diabetes Res* 1996;31:33-39.
33. Matsuba KT, Van Eeden SF, Saito Y, Okazawa M, Klut ME, Hayashi S, Hogg JC. Functional changes in aging polymorphonuclear leukocytes. *Circulation* 1998;97:91-98.
34. Wierusz-Wysocka B, Wykrętowicz A, Byks H, Sadurska K, Wysocki H. Polymorphonuclear neutrophils adherence, superoxide anion production (O²⁻) and HbA1 level in diabetic patients. *Diabetes Res Clin Pract* 1993;21:109-114.
35. Ceriello A, Bartolotti N, Motz E, Pieri C, Marra M, Tonutti L, Lizzio S, Felletto F, Catone B, Taboga C. Meal induced oxidative stress and low-density lipoprotein (LDL) oxidation in diabetes: the possible role of hyperglycemia. *Metabolism* 1999;48:1503-1508.
36. Mohanty P, Ghanim H, Hamouda W, Aljada A, Garg R, Dandona P. Both lipid and protein intakes stimulate increased generation of reactive oxygen species by polymorphonuclear leukocytes and mononuclear cells. *Am J Clin Nutr* 2002;5:767-772.
37. Brownlee M. The pathobiology of diabetic complications. *Unifying Mechanism Glycation and diabetic complications*. *Diabetes* 2005;54:1615-1625.